

**COVID 19 INFECTION AND VACCINATION:  
For breastfeeding, pregnant, and fertility-conscious women  
Carolyn P. Plourde, MD, FACOG**

1. Your risk of getting infected with COVID 19 is increased in pregnancy.  
TRUE FALSE
2. You are 2-4 times as likely to be admitted to an ICU, be intubated or die from COVID 19 infection during pregnancy.  
TRUE FALSE
3. Pregnant women were not included in the original vaccine trials.  
TRUE FALSE
4. The COVID vaccines have been shown to cause a slight reduction in fertility and increase risk of miscarriage.  
TRUE FALSE
5. Women who choose to be vaccinated against COVID 19 should try to receive the vaccine in the second trimester to avoid causing fetal birth defects (first trimester) and to keep the live virus from affecting the fetus (third trimester).  
TRUE FALSE
6. Women should wait six weeks after receiving the vaccine before trying to get pregnant.  
TRUE FALSE
7. A vaccinated mother transfers antibodies against COVID 19 through the placenta and through breastmilk.  
TRUE FALSE
8. Pregnant women have more side effects from the vaccine compared to non-pregnant women.  
TRUE FALSE
9. ACOG and SMFM are the two organizations that set the standards for obstetrical care in the United States.  
TRUE FALSE
10. Only the Moderna and Pfizer vaccines are safe in pregnancy; the Johnson & Johnson vaccine is to be avoided in pregnancy.  
TRUE FALSE